



Province of the  
**EASTERN CAPE**  
EDUCATION

Iphondo leMpuma Kapa: Isebe leMfundo  
Provinsie van die Oos Kaap: Departement van Onderwys  
Porafensie Ya Kapa Botjhabela: Lefapha la Thuto

# **NATIONAL SENIOR CERTIFICATE**

**GRADE 12**

**SEPTEMBER 2025**

**PHYSICAL SCIENCES P2: (CHEMISTRY)**

**MARKS: 150**

**TIME: 3 hours**

---

This question paper consists of 20 pages, including 4 data sheets.

---

**INSTRUCTIONS AND INFORMATION**

1. Write your name and surname in the appropriate space on the ANSWER BOOK.
2. This question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two sub questions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. Show ALL formulae and substitutions in ALL calculations.
9. Round off your FINAL numerical answers to a minimum of TWO decimal places.
10. Give brief motivations, discussions, etc. where required.
11. You are advised to use the attached DATA SHEETS.
12. Write neatly and legibly.

**QUESTION 1: MULTIPLE-CHOICE QUESTIONS**

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, for example 1.11 D.

- 1.1 Which ONE of the following homologous series can be described by the general formula  $C_nH_{2n}O$ ? (2)
- A Alcohols
- B Aldehydes
- C Esters
- D Carboxylic acids
- 1.2 The empirical formula of methyl propanoate is ... (2)
- A  $C_4H_8O_2$
- B  $C_4H_8O$
- C  $C_2H_4O$
- D  $CH_2O$
- 1.3 Consider the following compounds from the SAME HOMOLOGOUS SERIES with different boiling points.


Compounds	Boiling point (°C)
$C_4H_8O$	63
$C_4H_8O$	75

Which ONE of the following combinations is CORRECT regarding the homologous series to which compounds belong and type of isomers they form?

	Homologous series	Type of isomers
A	Ketones	Positional isomers
B	Ketones	Chain isomers
C	Aldehydes	Positional isomers
D	Aldehydes	Chain isomers


(2)

- 1.4 The organic molecule shown below undergoes thermal cracking.



Which ONE of the following are possible products from the reaction?


- A Hexane and butane
- B Octane, ethane and hydrogen gas
- C Pentane, propene and ethene
- D Propane, butane and ethene (2)
- 1.5 Which ONE of the following is TRUE regarding an EXOTHERMIC REACTION?
- A No energy is absorbed or released.
- B The amount of energy released is greater than the amount of energy absorbed.
- C The amount of energy absorbed is greater than the amount of energy released.
- D The amount of energy absorbed is equal to the amount of energy released. (2)
- 1.6 Consider the reaction below that reaches chemical equilibrium in a closed container.



Which ONE of the following changes will increase the yield of  $\text{NH}_3$ ?

- A Addition of a catalyst
- B Increase temperature
- C Decrease in the volume of the container
- D Decrease in the concentration of  $\text{N}_2$  (2)

1.7 The diagrams shows a titration set-up and titration curve obtained.



Consider the statements regarding substance **X**.

- I** The  $K_a$  value for substance **X** is greater than 1.
- II** The amount of **X** is equal the amount of NaOH at the equivalence point.
- III** Substance **X** ionises in water to produce low concentration of  $H_3O^+$

Which of the above statement(s) is/are TRUE?

- A **I** and **II** only
- B **II** only
- C **II** and **III** only
- D **I** and **III** only (2)

1.8 The reaction represented by the balanced equation below reaches equilibrium in a closed container.



Which ONE of the following changes will favour the reverse reaction?

- A Addition of hydrogen ( $H_2$ )
- B Addition of potassium hydroxide (KOH)
- C Decrease the pH of the equilibrium mixture
- D Cool the equilibrium mixture (2)


- 1.9 The standard hydrogen half-cell ( $\text{H}^+ | \text{H}_2 | \text{Pt}$ ) is connected to  $\text{Mn} | \text{Mn}^{2+}$  half-cell to form a galvanic cell.

Which ONE of the following combinations CORRECTLY identifies the oxidising agent and direction of electron flow?

	Oxidising agent	Direction of flow of electrons	
		From half-cell	To half-cell
A	Mn	$\text{H}^+   \text{H}_2   \text{Pt}$	$\text{Mn}   \text{Mn}^{2+}$
B	$\text{Mn}^{2+}$	$\text{Mn}   \text{Mn}^{2+}$	$\text{H}^+   \text{H}_2   \text{Pt}$
C	$\text{H}^+$	$\text{H}^+   \text{H}_2   \text{Pt}$	$\text{Mn}   \text{Mn}^{2+}$
D	$\text{H}^+$	$\text{Mn}   \text{Mn}^{2+}$	$\text{H}^+   \text{H}_2   \text{Pt}$

(2)

- 1.10 The electrolytic cell below is used for the electrolysis of concentrated sodium chloride ( $\text{NaCl}$ ).



Consider the statements regarding the electrolysis of concentrated sodium chloride ( $\text{NaCl}$ ).

- I** The pH of the electrolyte increases over time.
- II**  $\text{H}_2\text{O}$  is oxidised at electrode **P**.
- III** Chlorine gas ( $\text{Cl}_2$ ) is produced at electrode **Q**.

Which of the above statement(s) is/are TRUE?

- A **I** and **II** only
- B **III** only
- C **II** and **III** only
- D **I** and **III** only

(2)

**[20]**


**QUESTION 2 (Start on a NEW page.)**

The table below shows organic molecules (**A–F**) from different homologous series.

<b>A</b> $  \begin{array}{ccccccc}  & & & & \text{CH}_3 & & \\  & & & &   & & \\  \text{CH}_2 & - & \text{CH} & - & \text{CH}_2 & - & \text{CH} \\    & &   & & & &   \\  \text{CH}_3 & & \text{Cl} & & & & \text{CH}_3  \end{array}  $	<b>B</b> <p>3,4-dimethylpentan-2-ol</p>
<b>C</b> <p><math>\text{CH}_3\text{CH}(\text{CH}_3)\text{CCCH}_3</math></p>	<b>D</b> $  \begin{array}{ccccccc}  & \text{H} & & \text{H} & & \text{H} & & \text{O} \\  &   & &   & &   & &    \\  \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - & \text{C} - \text{H} \\  &   & &   & &   & & \\  & \text{H} & & & & \text{H} & & \\  & & &   & & & & \\  & & & \text{H} - \text{C} - \text{H} & & & & \\  & & &   & & & & \\  & & & \text{H} & & & &   \end{array}  $
<b>E</b> 3-methylbutanone	<b>F</b> $\text{C}_x\text{H}_y$

- 2.1 Define *homologous series*. (2)
- 2.2 Write down the LETTER that represents EACH of the following:
- 2.2.1 Unsaturated hydrocarbon (1)
- 2.2.2 A compound with a carbonyl group bonded to two saturated carbon atoms (1)
- 2.2.3 Compounds that are functional isomers (2)
- 2.3 Write down the IUPAC name of:
- 2.3.1 Compound **A** (3)
- 2.3.2 Compound **C** (2)
- 2.4 Draw the STRUCTURAL FORMULA of compound **B**. (2)
- 2.5 Compound **D** has a positional isomer.
- 2.5.1 Define the term *positional isomer*. (2)
- 2.5.2 Draw the positional isomer of compound **D**. (2)

- 2.6 Compound **F** ( $C_xH_y$ ) is an alkane that undergoes complete combustion with excess oxygen as shown below.



- 2.6.1 Determine the value **P**. (2)


- 2.6.2 Write down the CONDENSED STRUCTURAL FORMULA of compound **F**. (2)

[21]



**QUESTION 3 (Start on a NEW page.)**

The vapour pressure against temperature curves are drawn for THREE STRAIGHT CHAIN ALKANES. The vapour pressure are measured in mmHg. The atmospheric pressure is 760 mmHg.

**GRAPH OF VAPOUR PRESSURE VERSUS TEMPERATURE**

3.1 Define *vapour pressure*. (2)

3.2 At what phase are the alkanes at room temperature?

Write down only GAS, LIQUID or SOLID.

Give a reason for the answer. (2)

3.3 How will the boiling points change if the atmospheric pressure decreases?

Choose from INCREASES, DECREASES or REMAIN THE SAME. (1)

3.4 What can be concluded from the curve regarding the relationship between intermolecular forces and vapour pressure? (2)

3.5 The table below shows the boiling point of propan-1-ol and propanoic acid.

	COMPOUND	BOILING POINT (°C)
A	Propan-1-ol	97
B	Propanoic acid	141


3.5.1 Define the term *boiling point*. (2)

3.5.2 Fully explain the difference in the boiling points. (4)

[13]

**QUESTION 4 (Start on a NEW page.)**

Consider the flow diagram below:



4.1 Consider reaction I.

Write down the:

4.1.1 Name of reaction (1)

4.1.2 STRUCTURAL FORMULA of the carboxylic acid needed (2)

4.2 Consider reaction II.

Write down the:

4.2.1 NAME or FORMULA of the inorganic reagent needed (1)

4.2.2 IUPAC name of unsaturated hydrocarbon P (2)

4.3 Consider reaction III.

4.3.1 Write down the name of the type of addition reaction. (1)

4.3.2 Is compound Q a PRIMARY, SECONDARY or TERTIARY HALOALKANE? Give a reason for the answer. (2)

4.4 Compound Q is mixed with concentrated sodium hydroxide (NaOH) and strongly heated.

Write down the:

4.4.1 Type of reaction that occurred (1)


4.4.2 Balanced equation by using STRUCTURAL FORMULAE for the organic compounds and write the structural formula for the major product only in the reaction. (6)

[16]


**QUESTION 5 (Start on a NEW page.)**

5.1 Define the term *reaction rate*. (2)

A group of learners want to determine the percentage purity of a sample that contains calcium carbonate ( $\text{CaCO}_3$ ). They react 2,5 g sample of impure calcium carbonate ( $\text{CaCO}_3$ ) with EXCESS dilute hydrochloric acid ( $\text{HCl}$ ) according to the balanced equation.



They measure the volume of carbon dioxide gas produced over time and obtained the graph below.



5.2 How will the learners know that the reaction has reached completion? (1)

5.3 Calculate the:

5.3.1 Average rate at which  $\text{CO}_2$  is produced in  $\text{cm}^3 \cdot \text{s}^{-1}$  (3)


5.3.2 Percentage purity of  $\text{CaCO}_3$  if the molar volume of  $\text{CO}_2$  is  $24\,000 \text{ cm}^3 \cdot \text{mol}^{-1}$  at  $25^\circ\text{C}$  (5)

5.4 The experiment is repeated by using powdered sample of calcium carbonate instead of chunked sample of calcium carbonate.

5.4.1 Will the reaction rate INCREASE, DECREASE or REMAIN THE SAME? (1)

5.4.2 Explain the answer to QUESTION 5.4.1 by referring to the collision theory. (3)

- 5.5 The graph below represents the Maxwell-Boltzmann distribution curve for  $\text{CO}_2$  (g) at  $25^\circ\text{C}$ .



5.5.1 What does the shaded area to the right of line **P** represents? (1)

5.5.2 How will an increase in temperature affect the shaded area?


Choose from INCREASES, DECREASES or NO EFFECT.

Give a reason for the answer.

(2)  
[18]

**QUESTION 6 (Start on a NEW page.)**

When copper(II)chloride is dissolved in water the following chemical equilibrium equation is obtained. This can be used to demonstrate Le Chatelier's principle.



6.1 State *Le Chatelier's principle*. (2)

6.2 State whether the solution will become **BLUE** or **GREEN** when the following changes are made.


The colour of the equilibrium mixture is **BLUE**.

6.2.1 The mixture is cooled. (1)

6.2.2 Addition of saturated sodium chloride solution. (1)

6.3 Fully explain the answer to QUESTION 6.2.2 by referring to Le Chatelier's principle. (3)

6.4 Calcium carbonate is allowed to decompose at 1 000 °C in a sealed container according to the balanced chemical equation below.



Initially 0,24 mol of calcium carbonate is placed in a 2 dm<sup>3</sup> container and heated. An equilibrium is established at 1 000 °C. At this temperature the equilibrium constant is 0,0385.

6.4.1 Will the K<sub>c</sub> value INCREASE, DECREASE or REMAIN THE SAME if the temperature at which calcium carbonate decomposes is INCREASED?

Fully explain the answer. (3)

6.4.2 Calculate the mass of unreacted calcium carbonate that will remain in the container when the system reaches equilibrium at 1 000 °C. (7)

**[17]**

**QUESTION 7 (Start on a NEW page.)**

7.1 Consider the reactions below and answer questions that follow.

REACTION	CHEMICAL EQUATIONS
<b>A</b>	$\text{HSO}_4^- + \text{H}_2\text{O} \rightarrow \text{SO}_4^{2-} + \text{H}_3\text{O}^+$
<b>B</b>	$\text{KOH} + \text{CH}_3\text{COOH} \rightarrow \text{CH}_3\text{COOK} + \text{H}_2\text{O}$

7.1.1 Define a base in term of the *Arrhenius theory*. (2)

From the reactions above:

7.1.2 Which ONE of the reactions (**A** or **B**) represents the Arrhenius model? (1)

7.1.3 Write down a conjugate base of  $\text{HSO}_4^-$  (1)


7.2 Consider reaction **B**.

7.2.1 Give a reason why  $\text{CH}_3\text{COOH}$  can be regarded as a weak acid. (2)

7.2.2 Will  $\text{CH}_3\text{COOK}$  be ACIDIC, NEUTRAL or ALKALINE solution in water? (1)

7.2.3 Explain the answer in QUESTION 7.2.2 by using the relevant equation. (3)

7.3 During a titration,  $12,5 \text{ cm}^3$  of sodium hydroxide ( $\text{NaOH}$ ) solution with a concentration of  $0,2 \text{ mol}\cdot\text{dm}^{-3}$  neutralises  $25 \text{ cm}^3$  of sulphuric acid ( $\text{H}_2\text{SO}_4$ ) solution, according to the following balanced chemical equation:




7.3.1 Write down the name of a suitable indicator for this neutralisation reaction. (1)

7.3.2 Calculate the pH of the  $\text{H}_2\text{SO}_4$  solution. (7)

**[18]**

**QUESTION 8 (Start on a NEW page.)**

- 8.1 A galvanic cell is set up under standard conditions using unknown metal **M** and silver as an electrode. The standard electrode potential for the unknown metal **M** is less than 0 V.




- 8.1.1 Write down the energy conversion in this cell. (2)
- 8.1.2 Which electrode is the anode in the cell (Ag or Metal **M**)?  
Explain the answer. (3)
- 8.2 The initial potential difference for this cell is 1,54 V.
- 8.2.1 Identify metal **M** by calculation. (5)
- 8.2.2 Write down the cell notation for the cell. (3)
- 8.3 How will the initial reading on the voltmeter change when the following changes are made to the cell above?  
Write down only INCREASES, DECREASES or REMAINS THE SAME
- 8.3.1 Surface area of the Ag electrode is increased. (1)
- 8.3.2 The concentration of  $\text{AgNO}_3$  solution is increased to  $2 \text{ mol} \cdot \text{dm}^{-3}$ . (1)

**[15]**

**QUESTION 9 (Start on a NEW page.)**

The electrolytic cell, represented in the diagram below, is set up to purify copper which contains zinc and silver impurities.



9.1 Define the term *electrolyte*. (2)

9.2 Is this reaction ENDOTHERMIC or EXOTHERMIC? (1)

9.3 Which of the electrodes (**X** or **Z**) is connected to the positive terminal of the battery? (1)

9.4 Write down the half reaction that takes place at electrode **Z**. (2)

9.5 During the electrolysis it is observed that zinc is also oxidised but not silver.

Give a reason for this observation by referring to the relative strengths of the reducing agents. (1)

9.6 An unknown solution ( $MCl_2$ ) was electrolysed using copper electrodes. After some time  $1,806 \times 10^{22}$  electrons were gained at the cathode while the cathode gained 0,6 g.

Determine, by calculations, the formula of an unknown solution. (5)  
[12]

**TOTAL: 150**



**NATIONAL SENIOR CERTIFICATE  
NASIONALE SENIOR SERTIFIKAAT**

**DATA FOR PHYSICAL SCIENCES GRADE 12  
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESTE WETENSKAPPE GRAAD 12  
VRAESTEL 2 (CHEMIE)**

**TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES**

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	$p^\theta$	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume teen STD</i>	$V_m$	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	$T^\theta$	$273 \text{ K}$
Charge on electron <i>Lading op elektron</i>	$e$	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro se konstante</i>	$N_A$	$6,02 \times 10^{23} \text{ mol}^{-1}$

**TABLE 2: FORMULAE/TABEL 2: FORMULES**

$n = \frac{m}{M} \text{ or/of}$ $n = \frac{N}{N_A} \text{ or/of}$ $n = \frac{V}{V_m}$	$c = \frac{n}{V} \text{ or/of } c = \frac{m}{MV}$ $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$ $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} \text{ at /by } 298\text{K}$
$E^\theta_{\text{cell}} = E^\theta_{\text{cathode}} - E^\theta_{\text{anode}} / E^\theta_{\text{sel}} = E^\theta_{\text{katode}} - E^\theta_{\text{anode}}$ $E^\theta_{\text{cell}} = E^\theta_{\text{reduction}} - E^\theta_{\text{oxidation}} / E^\theta_{\text{sel}} = E^\theta_{\text{reduksie}} - E^\theta_{\text{oksidasie}}$ $E^\theta_{\text{cell}} = E^\theta_{\text{oxidising agent}} - E^\theta_{\text{reducing agent}} / E^\theta_{\text{sel}} = E^\theta_{\text{oksideermiddel}} - E^\theta_{\text{reduseermiddel}}$		
$q = I\Delta t \quad n = \frac{Q}{e} \quad \text{or/of } n = \frac{Q}{q_e}$		

TABLE 3: THE PERIODIC TABLE OF ELEMENTS/TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

1 (I)	2 (II)	3	4	5	6	7	8 Atoomgetal Atomic number	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	18 (VIII)
1 H 1							29 Cu 1,9										2 He 4
3 Li 7	4 Be 9						Elektronegatiwiteit Electronegativity	Simbool Symbol				5 B 11	6 C 12	7 N 14	8 O 16	9 F 19	10 Ne 20
11 Na 23	12 Mg 24						Approximate relative atomic mass/ Benaderde relatiewe atoommassa					13 Al 27	14 Si 28	15 P 31	16 S 32	17 Cl 35,5	18 Ar 40
19 K 39	20 Ca 40	21 Sc 45	22 Ti 48	23 V 51	24 Cr 52	25 Mn 55	26 Fe 56	27 Co 59	28 Ni 59	29 Cu 63,5	30 Zn 65	31 Ga 70	32 Ge 73	33 As 75	34 Se 79	35 Br 80	36 Kr 84
37 Rb 86	38 Sr 88	39 Y 89	40 Zr 91	41 Nb 92	42 Mo 96	43 Tc 98	44 Ru 101	45 Rh 103	46 Pd 106	47 Ag 108	48 Cd 112	49 In 115	50 Sn 119	51 Sb 122	52 Te 128	53 I 127	54 Xe 131
55 Cs 133	56 Ba 137	57 La 139	72 Hf 179	73 Ta 181	74 W 184	75 Re 186	76 Os 190	77 Ir 192	78 Pt 195	79 Au 197	80 Hg 201	81 Tl 204	82 Pb 207	83 Bi 209	84 Po 209	85 At 210	86 Rn 222
87 Fr 223	88 Ra 226	89 Ac															
			58 Ce 140	59 Pr 141	60 Nd 144	61 Pm 147	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175	
			90 Th 232	91 Pa 231	92 U 238	93 Np 237	94 Pu 244	95 Am 243	96 Cm 247	97 Bk 247	98 Cf 251	99 Es 252	100 Fm 257	101 Md 288	102 No 289	103 Lr 260	

TABLE 4A: STANDARD REDUCTION POTENTIALS  
TABEL 4A: STANDAARD REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies			$E^{\theta}$ (V)
$F_2(g) + 2e^-$	$\rightleftharpoons$	$2F^-$	+ 2,87
$Co^{3+} + e^-$	$\rightleftharpoons$	$Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^-$	$\rightleftharpoons$	$2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^-$	$\rightleftharpoons$	$Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^-$	$\rightleftharpoons$	$2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	$\rightleftharpoons$	$2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^-$	$\rightleftharpoons$	$2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^-$	$\rightleftharpoons$	$Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^-$	$\rightleftharpoons$	$Pt$	+ 1,20
$Br_2(l) + 2e^-$	$\rightleftharpoons$	$2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^-$	$\rightleftharpoons$	$NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^-$	$\rightleftharpoons$	$Hg(l)$	+ 0,85
$Ag^+ + e^-$	$\rightleftharpoons$	$Ag$	+ 0,80
$NO_3^- + 2H^+ + e^-$	$\rightleftharpoons$	$NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^-$	$\rightleftharpoons$	$Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^-$	$\rightleftharpoons$	$H_2O_2$	+ 0,68
$I_2 + 2e^-$	$\rightleftharpoons$	$2I^-$	+ 0,54
$Cu^+ + e^-$	$\rightleftharpoons$	$Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^-$	$\rightleftharpoons$	$S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^-$	$\rightleftharpoons$	$4OH^-$	+ 0,40
$Cu^{2+} + 2e^-$	$\rightleftharpoons$	$Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^-$	$\rightleftharpoons$	$SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^-$	$\rightleftharpoons$	$Cu^+$	+ 0,16
$Sn^{4+} + 2e^-$	$\rightleftharpoons$	$Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^-$	$\rightleftharpoons$	$H_2S(g)$	+ 0,14
<b><math>2H^+ + 2e^-</math></b>	<b><math>\rightleftharpoons</math></b>	<b><math>H_2(g)</math></b>	<b>0,00</b>
$Fe^{3+} + 3e^-$	$\rightleftharpoons$	$Fe$	- 0,06
$Pb^{2+} + 2e^-$	$\rightleftharpoons$	$Pb$	- 0,13
$Sn^{2+} + 2e^-$	$\rightleftharpoons$	$Sn$	- 0,14
$Ni^{2+} + 2e^-$	$\rightleftharpoons$	$Ni$	- 0,27
$Co^{2+} + 2e^-$	$\rightleftharpoons$	$Co$	- 0,28
$Cd^{2+} + 2e^-$	$\rightleftharpoons$	$Cd$	- 0,40
$Cr^{3+} + e^-$	$\rightleftharpoons$	$Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^-$	$\rightleftharpoons$	$Fe$	- 0,44
$Cr^{3+} + 3e^-$	$\rightleftharpoons$	$Cr$	- 0,74
$Zn^{2+} + 2e^-$	$\rightleftharpoons$	$Zn$	- 0,76
$2H_2O + 2e^-$	$\rightleftharpoons$	$H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^-$	$\rightleftharpoons$	$Cr$	- 0,91
$Mn^{2+} + 2e^-$	$\rightleftharpoons$	$Mn$	- 1,18
$Al^{3+} + 3e^-$	$\rightleftharpoons$	$Al$	- 1,66
$Mg^{2+} + 2e^-$	$\rightleftharpoons$	$Mg$	- 2,36
$Na^+ + e^-$	$\rightleftharpoons$	$Na$	- 2,71
$Ca^{2+} + 2e^-$	$\rightleftharpoons$	$Ca$	- 2,87
$Sr^{2+} + 2e^-$	$\rightleftharpoons$	$Sr$	- 2,89
$Ba^{2+} + 2e^-$	$\rightleftharpoons$	$Ba$	- 2,90
$Cs^+ + e^-$	$\rightleftharpoons$	$Cs$	- 2,92
$K^+ + e^-$	$\rightleftharpoons$	$K$	- 2,93
$Li^+ + e^-$	$\rightleftharpoons$	$Li$	- 3,05

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë

**TABLE 4B: STANDARD REDUCTION POTENTIALS**  
**TABEL 4B: STANDAARD REDUKSIEPOTENSIALE**

Half-reactions/Halfreaksies		$E^{\theta}$ (V)
$\text{Li}^{+} + \text{e}^{-}$	$\rightleftharpoons$ Li	- 3,05
$\text{K}^{+} + \text{e}^{-}$	$\rightleftharpoons$ K	- 2,93
$\text{Cs}^{+} + \text{e}^{-}$	$\rightleftharpoons$ Cs	- 2,92
$\text{Ba}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Ba	- 2,90
$\text{Sr}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Sr	- 2,89
$\text{Ca}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Ca	- 2,87
$\text{Na}^{+} + \text{e}^{-}$	$\rightleftharpoons$ Na	- 2,71
$\text{Mg}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Mg	- 2,36
$\text{Al}^{3+} + 3\text{e}^{-}$	$\rightleftharpoons$ Al	- 1,66
$\text{Mn}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Mn	- 1,18
$\text{Cr}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Cr	- 0,91
$2\text{H}_2\text{O} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{H}_2(\text{g}) + 2\text{OH}^{-}$	- 0,83
$\text{Zn}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Zn	- 0,76
$\text{Cr}^{3+} + 3\text{e}^{-}$	$\rightleftharpoons$ Cr	- 0,74
$\text{Fe}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Fe	- 0,44
$\text{Cr}^{3+} + \text{e}^{-}$	$\rightleftharpoons$ $\text{Cr}^{2+}$	- 0,41
$\text{Cd}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Cd	- 0,40
$\text{Co}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Co	- 0,28
$\text{Ni}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Ni	- 0,27
$\text{Sn}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Sn	- 0,14
$\text{Pb}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Pb	- 0,13
$\text{Fe}^{3+} + 3\text{e}^{-}$	$\rightleftharpoons$ Fe	- 0,06
$2\text{H}^{+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{H}_2(\text{g})$	<b>0,00</b>
$\text{S} + 2\text{H}^{+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{H}_2\text{S}(\text{g})$	+ 0,14
$\text{Sn}^{4+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{Sn}^{2+}$	+ 0,15
$\text{Cu}^{2+} + \text{e}^{-}$	$\rightleftharpoons$ $\text{Cu}^{+}$	+ 0,16
$\text{SO}_4^{2-} + 4\text{H}^{+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+ 0,17
$\text{Cu}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Cu	+ 0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^{-}$	$\rightleftharpoons$ $4\text{OH}^{-}$	+ 0,40
$\text{SO}_2 + 4\text{H}^{+} + 4\text{e}^{-}$	$\rightleftharpoons$ $\text{S} + 2\text{H}_2\text{O}$	+ 0,45
$\text{Cu}^{+} + \text{e}^{-}$	$\rightleftharpoons$ Cu	+ 0,52
$\text{I}_2 + 2\text{e}^{-}$	$\rightleftharpoons$ $2\text{I}^{-}$	+ 0,54
$\text{O}_2(\text{g}) + 2\text{H}^{+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{H}_2\text{O}_2$	+ 0,68
$\text{Fe}^{3+} + \text{e}^{-}$	$\rightleftharpoons$ $\text{Fe}^{2+}$	+ 0,77
$\text{NO}_3^{-} + 2\text{H}^{+} + \text{e}^{-}$	$\rightleftharpoons$ $\text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+ 0,80
$\text{Ag}^{+} + \text{e}^{-}$	$\rightleftharpoons$ Ag	+ 0,80
$\text{Hg}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{Hg}(\ell)$	+ 0,85
$\text{NO}_3^{-} + 4\text{H}^{+} + 3\text{e}^{-}$	$\rightleftharpoons$ $\text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+ 0,96
$\text{Br}_2(\ell) + 2\text{e}^{-}$	$\rightleftharpoons$ $2\text{Br}^{-}$	+ 1,07
$\text{Pt}^{2+} + 2\text{e}^{-}$	$\rightleftharpoons$ Pt	+ 1,20
$\text{MnO}_2 + 4\text{H}^{+} + 2\text{e}^{-}$	$\rightleftharpoons$ $\text{Mn}^{2+} + 2\text{H}_2\text{O}$	+ 1,23
$\text{O}_2(\text{g}) + 4\text{H}^{+} + 4\text{e}^{-}$	$\rightleftharpoons$ $2\text{H}_2\text{O}$	+ 1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^{+} + 6\text{e}^{-}$	$\rightleftharpoons$ $2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+ 1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^{-}$	$\rightleftharpoons$ $2\text{Cl}^{-}$	+ 1,36
$\text{MnO}_4^{-} + 8\text{H}^{+} + 5\text{e}^{-}$	$\rightleftharpoons$ $\text{Mn}^{2+} + 4\text{H}_2\text{O}$	+ 1,51
$\text{H}_2\text{O}_2 + 2\text{H}^{+} + 2\text{e}^{-}$	$\rightleftharpoons$ $2\text{H}_2\text{O}$	+ 1,77
$\text{Co}^{3+} + \text{e}^{-}$	$\rightleftharpoons$ $\text{Co}^{2+}$	+ 1,81
$\text{F}_2(\text{g}) + 2\text{e}^{-}$	$\rightleftharpoons$ $2\text{F}^{-}$	+ 2,87

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë